

Fertilization of Strawberries in Florida¹

George Hochmuth and Earl Albregts²

INTRODUCTION

Strawberries are produced on 5000-6000 acres each year in Florida and the 1992-93 season had a crop value of \$103 million (Freie and Pugh, 1994). Although the major strawberry production area is in Hillsborough county, there are numerous other production areas throughout the state, notably Alachua, Bradford, Dade, Manatee, Palm Beach, and Polk counties (Hochmuth, 1988).

Production costs for strawberries average \$14,000 per acre; however, fertilizer accounts for only 2% of the total costs (Smith and Taylor, 1993). Although fertilizer comprises only a small fraction of strawberry production costs, proper fertilization is important for maximizing yield and fruit quality, and for minimizing potential negative impact to the environment caused by leaching or runoff of excess fertilizer.

FERTILIZATION RECOMMENDATIONS

Strawberry P and K fertilization recommendations are based on the calibrated Mehlich-1 soil test results and vary according to the

soil-test level of P and K (Table 1). N requirements are 150 lb N per acre for the total season (200 days) (Hochmuth and Hanlon (in press)). Growers should avoid the temptation to apply more N because excess N can increase the amount of malformed fruits (Albregts and Howard, 1982) and can lead to excessive vegetative plant growth and reduced yields. Rates of K higher than recommended reduced average strawberry fruit size and yield (Albregts et al., 1991; Albregts et al., 1994).

DRIP IRRIGATION

Drip irrigation can result in up to 50% reduction in water use for strawberries, compared to overhead sprinklers (Myers and Locascio, 1972). Nearly all strawberries in Florida now are irrigated with drip irrigation systems. Although drip irrigation can improve irrigation efficiency, care must be exercised to operate the system properly so optimum amounts of water are applied. Inadequate irrigation can reduce yields and over-irrigation can leach N and K. Irrigation amounts should be scheduled to meet the strawberry crop evapotranspiration needs which range from 800 gal per acre per day in October to 3000 gal per acre per day in April (Clark, 1993).

1. This document is CIR1141, one of a series of the Horticultural Sciences Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida. Original publication date September 1994. Reviewed May 2003. Visit the EDIS Web Site at <http://edis.ifas.ufl.edu>.

2. George Hochmuth, Professor, Horticultural Sciences Department and Earl Albregts, Professor, Agricultural Research and Educational Center, Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida, Gainesville FL 32611.

FERTIGATION

Applying fertilizer through the drip irrigation system (fertigation) offers the potential for increasing efficiency of application of leachable nutrients such as N and K (Locascio and Myers, 1975; Locascio and Martin, 1985). With fertigation, N and K can be applied in small amounts corresponding to crop N and K requirements. Current recommendations call for incorporating all the P and micronutrients and up to 30 lb per acre each N and K₂O in the beds preplant. (Plate 1) The remaining N and K can be injected through the drip irrigation system.



(Plate 1).

N and K can be injected on a daily or weekly basis. Schedules for N and K injection were presented for strawberries (Hochmuth and Clark, 1991); however, recent research has made it possible to reduce strawberry fertilization rates (Hochmuth et al., 1994; Albrechts et al., 1994). Fertilization rates can be reduced from 1.0 lb N and K₂O each per acre per day. Revised fertigation recommendations are presented in Table 2 .

PETIOLE SAP-TESTING


Timely plant tissue analysis for N and K is important for optimizing N and K injection and for aid in making decisions regarding amounts of N and K to inject. Using research data from Dover AREC and commercial strawberry farms, extension scientists developed petiole fresh sap-testing procedures. The petiole sap sufficiency ranges for N and K are presented in Table 3. There are several sap-testing kits available for determining sap N and K concentrations. (Plate 2) The brochure describing the procedures available from the county Extension offices, is entitled "Guide for Plant Petiole Sap-Testing for Vegetable Crops."



(Plate 2).

WHOLE-LEAF TESTING

Petiole sap analyses should be backed up by routine analyses of whole leaves by a tissue testing laboratory. (Plate 3) Presently, sap testing is limited to N and K because either test kits do not exist or procedures have not been developed for other nutrients. Sufficiency ranges for nutrients in whole strawberry leaves are presented in Table 4.



(Plate 3).

CHECKLIST FOR SUCCESSFUL STRAWBERRY DRIP FERTIGATION

- Practice calibrated soil testing to determine lime and fertilizer requirements.
- Select correct rates of fertilizer to apply to soil before bedding and mulching. Preplant (broadcast) N and K₂O should not exceed 30 lb each per acre.
- Practice careful irrigation management to avoid over-irrigation which leaches N and K from the root zone.

- Inject N and K in daily or weekly injections at rates not exceeding 0.60 lb each N and K₂O per acre per day except for 0.75 lb each N and K₂O per acre per day in February and March.
- Monitor petiole sap N and K concentrations and use that information to adjust N and K injection schedules or rates.
- Back up petiole sap-testing with laboratory analyses of whole leaves.

LITERATURE

- Albregts, E. E., G. J. Hochmuth, C. K. Chandler, and J. A. Cornell. 1994. K requirements for drip irrigated strawberry. Unpublished data.
- Albregts, E. E., and C. M. Howard. 1982. Effect of fertilizer rate on number of malformed strawberry fruit. *Proc. Fla. State Hort. Soc.* 95:323-324.
- Albregts, E. E., C. M. Howard, and C. K. Chandler. 1991. Strawberry responses to K rate on a fine sand soil. *HortScience* 26:135-138.
- Clark, G. A. 1993. Water requirements for drip-irrigated strawberries in south central Florida. *Gulf Coast Res. and Ed. Center Res. Report BRA* 1993-2.
- Freie, R. L., and N. L. Pugh. 1994. Florida Agricultural Statistics. Vegetable Summary. Fla. Dept. Agric. Consumer Serv. Tallahassee, FL.
- Hochmuth, G. J., ed. 1988. Strawberry production guide for Florida. Fla. Coop. Ext. Serv. Circ. 142C.
- Hochmuth, G. J., and G. A. Clark. 1991. Fertilizer application and management for micro- (or drip) irrigated vegetables in Florida. Fla. Coop. Ext. Serv. Special Series SS-VEC-45.
- Hochmuth, G. J., and E. A. Hanlon. In Press. Commercial vegetable crop nutrient requirements. Fla. Coop. Ext. Serv. Circ. 806.
- Hochmuth, G. J., D. N. Maynard, C. S. Vavrina, and E. A. Hanlon. 1991. Plant tissue analysis and interpretation for vegetable crops in Florida. Fla. Coop. Ext. Serv. Special Series SS-VEC-42.
- Hochmuth, G. J., E. E. Albregts, C. K. Chandler, and J. A. Cornell. 1994. N requirements for drip-irrigated strawberry. Unpublished data.
- Kidder, G., E. A. Hanlon, and G. J. Hochmuth. 1989. IFAS standardized fertilization recommendations for vegetable crops. Fla. Coop. Ext. Serv. Special Series SS-SOS-907.
- Locascio, S. J., and J. M. Myers. 1975. Trickle irrigation and fertilization method for strawberries. *Proc. Fla. State Hort. Soc.* 88:185-189.
- Locascio, S. J., and F. G. Martin. 1985. Nitrogen source and application timing for trickle-irrigated strawberries. *J. Amer. Soc. Hort. Sci.* 110:820-823.
- Myers, J. M., and S. J. Locascio. 1972. Efficiency of irrigation methods for strawberries. *Proc. Fla. State Hort. Soc.* 85:114-117.
- Smith, S. A., and T. G. Taylor. 1993. Production cost for selected vegetables in Florida, 1992-1993. Fla. Coop. Ext. Serv. Circ. 1121.

Table 1.

Table 1. Interpretation of Mehlich-1 soil test and revised nutrient recommendations for strawberries.					
Element	Very Low	Low	Medium	High	Very High
	parts per million soil				
P	<10	10-15	16-30	31-60	>60
K	<20	20-35	36-60	61-125	>125
Fertilizer	lb per acre				
P ₂ O ₅	150	120	100	0	0
K ₂ O	150	120	100	0	0

Table 2.

Table 2. N and K fertigation recommendations for strawberries.		
Stage in season	Injection rate (lb/acre/day) ^z	
	N	K ₂ O
First 2 weeks	0.3	0.3
February/ March	0.75	0.75
All other months	0.6	0.6
^z Based on cultural system with beds on four-ft centers and up to 30 lb each N and K and K ₂ O per acre broadcast in bed before planting on soils testing very low in K.		
K injection rates drop by about 20% as soil test goes from VL to L and again from L to Medium.		

Table 3.

Table 3. Sufficiency ranges for petiole sap N and K concentrations for strawberries, October planting.		
Month of season	Petiole sap nutrient concentration(ppm)	
	NO ₃ - N	K
November	800 - 900	3000 - 3500
December	600 - 800	3000 - 3500
January	600 - 800	2500 - 3000
February	300 - 500	2000 - 2500
March	200 - 500	1800 - 2500
April	200 - 500	1500 - 2000

Table 4.

Table 4. Sufficiency nutrient ranges in most-recently-matured whole leaves (blade plus petioles) of strawberries													
Time of Sampling	Status	N	P	K	Ca	Mg	S	Fe	Mn	Zn	B	Cu	Mo
		ppm											
		%											
Initial Harvest	Deficient	<3.0	0.2	1.5	0.4	0.25	0.25	50	30	20	20	5	-- ^z
	Adequate Range	3.0 -3.5	0.2 -0.4	1.5 -2.5	0.4 -1.5	0.25 -0.50	0.25 -0.80	50 -100	30 -100	20 -40	20 -40	5 -10	--
	High	>3.5	0.4	2.5	1.5	0.50	0.80	100	100	40	40	10	--
	Toxic (>)	--	--	--	--	--	--	--	800	--	--	--	--
Midseason	Deficient	<2.8	0.2	1.1	0.4	0.2	0.25	50	25	20	20	5	0.5
	Adequate Range	2.8 -3.0	0.2 -0.4	1.1 -2.5	0.4 -1.5	0.2 -0.4	0.25 -0.80	50 -100	25-100	20 -40	20 -40	5 -10	0.5-0.8
	High	>3.0	0.4	2.5	1.5	0.4	0.80	100	100	40	40	10	0.8
	Toxic (>)	--	--	--	--	--	--	--	800	--	--	-	--
^z Values not available													